

File format for GONG solar model project.

Jørgen Christensen-Dalsgaard

Stellar Astrophysics Centre; and Department of Physics and Astronomy, Aarhus University

8 October, 2015

1. Introduction.

The present notes provide a definition of the data format used for comparison of solar models within the GONG models team. They essentially correspond to Section 9 of *Computational procedures for GONG model project* (in the following **CP**) and have been extracted here for convenience of reference. For details on the GONG model comparisons project, involving simplified physics, **CP** should be consulted. The focus here is on the exchange of realistic solar models, involving detailed physics.

The use of the format has been extended to more general exchange of stellar models, including in model comparisons related to the *Kepler* project.

The main update in the present version is the introduction of a new data format, corresponding essentially to full binary double precision. This is discussed in Section 2.4.

2. File structure, and variables

The exchange of models will take place by means of ASCII files, to avoid problems with different binary formats. This makes it possible to send the data electronically without problems (other than caused by the amount).

For each model the file consists of a header, with descriptive information, a set of global variables, and a set of variables given at each mesh point.

The set of variables is likely to develop, particularly in the context of the use of the format for transferring realistic solar and stellar models. The actual set used is defined by the variable `ivers` included in the header to the file; this includes `ivers0` depending on the variable set, and a flag for the output format (see also Section 2.4). In addition, the total number of global parameters, meshpoints and variables at each meshpoint are given in the header, so that the information required to read the file is available.

The variables should be given in *cgs* units, unless otherwise stated.

2.1 Text header.

The first record should contain the name of the model, its date, and an identification of its origin (such as name or institute). In addition the header may contain text further describing the calculation, and information about the remaining data.

An example of a header may be the following (describing Model S of the *Science* series of articles):

```
L5BI.D.15.PRES.950912.AARHUS
Level 5 physics, present Sun. (OPAL opacity, OPAL EOS). He, Z diffusion.
Age of present Sun: 4.6 Gyr.
```

2.2 Global parameters.

These are set up in the array `glob(i)`, $i = 1, \dots, \text{iconst}$, with the following definition:

- 1: M (total mass).
- 2: R (photospheric radius).
- 3: L_s (surface luminosity).
- 4: Z_0 (initial heavy-element abundance).^{a)}
- 5: X_0 (initial hydrogen abundance).
- 6: $\alpha = \ell/H_p$ (mixing-length parameter; H_p is pressure scale height)
- 7: ϕ (another convection theory parameter)^{b)}
- 8: ξ (yet another convection-theory parameter)^{b)}
- 9: β (parameter in surface-pressure condition)^{c)}
- 10: λ (parameter in surface-luminosity condition)^{c)}
- 11: $\frac{R^2}{p_c} \frac{d^2 p_c}{dr^2}$ (at centre)^{d)}
- 12: $\frac{R^2}{\rho_d} \frac{d^2 \rho_c}{dr^2}$ (at centre)^{d)}
- 13: Model age (in years)^{e)}
- 14: T_{eff} (effective temperature, in K)^{f)}
- 15: Gravitational constant G , in cgs units^{g)}

Notes:

- a) Text changed 18/2/13 to emphasize that this should be the *initial* heavy-element abundance.
- b) The parameters ϕ and ξ are defined in **CP**. For the Böhm-Vitense formulation, $\phi = 9/4$ and $\xi = 1/162$.
- c) These parameters are used in boundary conditions for models with simplified physics (*cf.* **CP**). For realistic models, $\beta = \lambda = 1$.
- d) These second derivatives may be needed in the central boundary conditions for oscillation calculations.
- e) Added 30/7/96. If the calculation includes pre-main-sequence evolution, the definition of zero age must be specified, in the character header or in accompanying notes. It would be useful to arrive at a common definition; suggestions are welcome.
It appears that this was not properly included (at least in some models) before 9/1/13.
- f) Added 9/1/13.
- g) Added 30/4/14.

2.3 Model variables at each mesh point.

These are set up in the array `var(i, n)`, $i = 1, \dots, \text{ivar}$, $n = 1, \dots, \text{nn}$. The current set of variables, as defined in October 2005 is characterized by having `ivers0 = 300`.

- 1: r (distance to centre)
- 2: $\ln q$, $q = m/M$ (m is mass interior to r and M is total mass)
- 3: T (temperature)
- 4: p (pressure)
- 5: ρ (density)
- 6: X (hydrogen abundance by mass)
- 7: $L(r)$ (luminosity at distance r from centre)

- 8: κ (opacity)
- 9: ϵ (energy generation rate per unit mass)
- 10: $\Gamma_1 = \left(\frac{\partial \ln p}{\partial \ln \rho} \right)_{\text{ad}}$
- 11: $\nabla_{\text{ad}} = \left(\frac{\partial \ln T}{\partial \ln p} \right)_{\text{ad}}$
- 12: $\delta = - \left(\frac{\partial \log \rho}{\partial \log T} \right)_p$
- 13: c_p (specific heat at constant pressure)
- 14: μ_e^{-1} [see note ii) below]
- 15: $\frac{1}{\Gamma_1} \frac{d \log p}{d \log r} - \frac{d \log \rho}{d \log r}$
- 16: r_X (rate of change in X from nuclear reactions)
- 17: Z (heavy-element abundance per unit mass)
- 18: $R - r$
- 19: ϵ_g (rate of gravitational energy release)
- 20: L_g (local gravitational luminosity; this has only been included in J. Reiter's models so far)
- 21: $X(^3\text{He})$ (^3He abundance by mass)
- 22: $X(^{12}\text{C})$ (^{12}C abundance by mass)
- 23: $X(^{13}\text{C})$ (^{13}C abundance by mass)
- 24: $X(^{14}\text{N})$ (^{14}N abundance by mass)
- 25: $X(^{16}\text{O})$ (^{16}O abundance by mass)
- 26: $\left(\frac{\partial \ln \Gamma_1}{\partial \ln \rho} \right)_{p,Y}$
- 27: $\left(\frac{\partial \ln \Gamma_1}{\partial \ln p} \right)_{\rho,Y}$
- 28: $\left(\frac{\partial \ln \Gamma_1}{\partial Y} \right)_{p,\rho}$
- 29: $X(^2\text{H})$ (^2H abundance by mass)
- 30: $X(^4\text{He})$ (^4He abundance by mass)
- 31: $X(^7\text{Li})$ (^7Li abundance by mass)
- 32: $X(^7\text{Be})$ (^7Be abundance by mass)
- 33: $X(^{15}\text{N})$ (^{15}N abundance by mass)
- 34: $X(^{17}\text{O})$ (^{17}O abundance by mass)
- 35: $X(^{18}\text{O})$ (^{18}O abundance by mass)
- 36: $X(^{20}\text{Ne})$ (^{20}Ne abundance by mass)
- 37 – 40: Currently not used.

Comments:

- i) In variable 2, $\ln q$ is used instead of m to give a better indication of variation close to the surface. Here “ln” is natural logarithm.

- ii) $\mu_e^{-1} = N_e m_u$, where N_e is the number of free electrons per unit mass and m_u is the atomic mass unit; thus μ_e is the mean molecular weight per electron. It has been included to give some indication of the ionization state.
- iii) Variable 18 was introduced Aug. 6 1993, to avoid problems with interpolation in r near surface. (For versions before `ivers0 = 210`, but after that date, $R - r$ was set in variable 17.)
- iv) The derivatives of Γ_1 were introduced principally to allow calculation of kernels involving Y , in realistic models; they are less essential for the comparison of models.
- v) Calculations that do not follow all details of the CNO cycle may not have available some of the CNO abundances. In that case, the corresponding abundances can be set to zero. (An example might be a calculation following the conversion of ^{16}O into ^{14}N but ignoring the details of carbon burning into ^{14}N .)
- vi) More generally, for a given code there might be variables that are difficult to obtain. In that case, the above numbering should be maintained, but the missing variables may be set identically to zero.

The present set of variables has been chosen to give a reasonably comprehensive basis for comparing evolution models, and to be adequate for the computation of adiabatic oscillations. In particular these variables should define completely the stellar structure equations, and so permit a check of the accuracy to which the equations are satisfied. For a more detailed comparison even more variables may be needed (such as ionization levels), but I suggest that that be arranged separately.

For non-adiabatic calculations, which we should eventually get to, more variables are certainly needed. Typical examples are the derivatives of κ and ϵ with respect to p and T , and possibly variables relating to the perturbation of the convective flux. They can be included later, by extending the basic set given above.

2.4 Format for data transfer.

The data exchange should be carried out by means of formatted ASCII files, using the following structure:

Record 1: Name of model (as a character string)
Record 2 – 4: Explanatory text, in free format
Record 5: `nn, iconst, ivar, ivers`
Record 6 – 8: `glob(i), i = 1, ..., iconst`
Record 9 – : `var(i,n), i = 1, ..., ivar, n = 1, ... nn.`

Here `nn` is the number of mesh points in the model, `iconst` is the number of global variables (given in the array `glob`), and `ivar` is the number of variables at each mesh point (given in the array `var`). Thus with the present set `iconst = 15` and `ivar = 40`, but there is room for expansion. The version number `ivers` is written as `ivers = ivers1 + ivers0`. Here `ivers0` was defined above; as discussed below `ivers1 = 0` corresponds to the old (before 8 October 2015) data format, while `ivers1 = 1000` should be used for the new more precise data format.

The integer variable Record 5 should be written with the format

```
4i10
```

There are two options for the format for the real-variable Records 6 and up. Before 8 October 2015 the only format was

```
1p5e16.9
```

The revision has implemented the format

1p,5(x,e26.18e3)

The use of this format must be flagged by adding 1000 in `ivers`.

3. Summary of earlier versions

The format has been in used for more than a decade, with various extensions or modifications; these are characterized by the version number, as summarized below:

- **Version 100:** `var(1)` - `var(16)` defined as above. In some models `var(17)` was set to $R - r$, and `var(19)` and `var(20)` were set as above; otherwise these variables were set to zero; `ivar` = 20.
- **Version 200:** `var(1)` - `var(16)` and `var(19)` - `var(25)` defined as above, `var(17)` set to $R - r$; `ivar` = 25.
- **Version 210:** `var(1)` - `var(25)` defined as above; `ivar` = 25.
- **Version 250:** `var(1)` - `var(28)` defined as above; `ivar` = 30.